

### Immunology Referral Form

Please complete the following and fax with clinical documentation to:  
844.797.5050 or email CenterSource@soleohealth.com

#### Referral Process

1. PATIENT INFORMATION			2. PHYSICIAN INFORMATION		
Name:			Physician's name:		
Address:			Address:		
City:	State:	Zip:	City:	State:	Zip:
Home Phone:		Other Phone:	Office Contact:		
Email:			Phone:		Fax:
DOB:		Social Security #:		NPI:	
Gender: <input type="checkbox"/> M <input type="checkbox"/> F		Height:	Weight:		
Allergies: _____					

#### 3. DIAGNOSIS Year of diagnosis: \_\_\_\_\_

<input type="checkbox"/> Hereditary Hypogammaglobulinemia (D80.0)
<input type="checkbox"/> Selective deficiency of immunoglobulin A (IgA) (D80.2)
<input type="checkbox"/> Selective deficiency of immunoglobulin G (IgG) subclasses (D80.3)
<input type="checkbox"/> Selective deficiency of immunoglobulin M (IgM) (D80.4)
<input type="checkbox"/> Immunodeficiency with Increased immunoglobulin M (IgM) (D80.5)
<input type="checkbox"/> Antibody deficiency with near-normal immunoglobulins or with hyperimmunoglobulinemia (D80.6)
<input type="checkbox"/> Transient hypogammaglobulinemia of infancy (80.7)
<input type="checkbox"/> Severe combined immunodeficiency (SCID) with reticular dysgenesis (D81.0)
<input type="checkbox"/> Severe combined immunodeficiency (SCID) with low T and B cell numbers (D81.1)
<input type="checkbox"/> Severe combined immunodeficiency (SCID) with low or normal B cell numbers (D81.2)
<input type="checkbox"/> Purine nucleoside phosphorylase (PNP) deficiency (D81.5)
<input type="checkbox"/> Major histocompatibility complex class I deficiency (D81.6)
<input type="checkbox"/> Major histocompatibility complex class II deficiency (D81.7)
<input type="checkbox"/> Other Combined Immunodeficiencies (D81.89)
<input type="checkbox"/> Combined Immunodeficiencies, unspecified (D81.9)
<input type="checkbox"/> Wiskott-Aldrich Syndrome (D82.0)
<input type="checkbox"/> Di George's syndrome (D82.1)
<input type="checkbox"/> Hyperimmunoglobulin E (IgE) syndrome (D82.4)
<input type="checkbox"/> Common variable immunodeficiency with predominant abnormalities of B-cell numbers (D83.0)
<input type="checkbox"/> Common variable immunodeficiency with predominant immunoregulatory T-cell disorders (D83.1)
<input type="checkbox"/> Common variable immunodeficiency with autoantibodies to B or T cells (D83.2)
<input type="checkbox"/> Other common variable immunodeficiencies (D83.8)
<input type="checkbox"/> Common variable immunodeficiency (CVID), unspecified (D83.9)
<input type="checkbox"/> Other Specified Immunodeficiencies (D84.8)
<input type="checkbox"/> Primary Immunodeficiencies (D84.9)
<input type="checkbox"/> Cerebellar ataxia with defective DNA repair (G11.3)
<input type="checkbox"/> Other(s):

#### 4. INSURANCE INFORMATION

Please submit copies of the front and back of primary and secondary insurance cards with this referral.

**5. ADDITIONAL INFORMATION REQUESTED**

Has the Patient Received IVIG Previously?  No  Yes Product: \_\_\_\_\_  
Date of last dose: \_\_\_\_\_  
Last BUN/CR \_\_\_\_\_  IgA level \_\_\_\_\_  H&P  Infection History  Baseline IgG level  
 Immune Response to Vaccines

**6. PRESCRIPTION INFORMATION** Anticipated Start Date: \_\_\_\_\_

**Immune Globulin** Product: \_\_\_\_\_  IV  SQ  
Administer \_\_\_\_\_ grams daily for \_\_\_\_\_ day(s) OR \_\_\_\_\_ milligrams/kilogram daily over \_\_\_\_\_ day(s)  
Repeat course every \_\_\_\_\_ week(s) for a total of \_\_\_\_\_ courses/cycles  
 Pre-hydrate with:  NS  D5W  Other: \_\_\_\_\_ ml IV over \_\_\_\_\_ hours  
 Pre medicate:  
 Acetaminophen 325-650mg PO prior to IG and then Q4-6 hours as needed during infusion  
 Diphenhydramine 25-50 mg PO prior to IG  
 Other premedication: \_\_\_\_\_  
 Post-hydrate with:  NS  D5W  Other: \_\_\_\_\_ ml IV over \_\_\_\_\_ hours  
Provide supplies necessary to maintain IV Access:  PIV  Midline/PICC  Port  
Administration method:  Pump  Dial-a-flow

**7. FLUSH ORDERS**

**PIV/midline/PICC:** Flush before, after each infusion, and as needed with 3-20 ml NS, followed by Heparin 2-5 ml if indicated  
 Heparin 10 unit/ml  Heparin 100 unit/ml  
**Port:** Flush before, after each infusion, and as needed with 5-20 ml NS, followed by Heparin 100 unit/ml 5 ml. (Use Heparin 10 unit/ml for patients <2 years old)

**8. ANAPHYLAXIS ORDERS**

**Adults:** For mild reaction-administer diphenhydramine 50mg IVP/PO x 1.  
For severe reactions-administer epinephrine 0.3mg SubQ/IM x 1. May repeat x1 if needed. Contact 911 and monitor vital signs.  
**Pediatrics:** administer by age: For mild reaction (rash/hives) give diphenhydramine  
**Age 1-5:** 12.5ml IV/PO x1 **Age 6-11:** 25mg IV/PO x1 **Age 12+:** 50mg IV/PO x1  
For severe reaction (airway closure) administer epinephrine. 0.01mg/kg/dose (max 0.3mg) SubQ/IM x 1.  
If needed, repeat a second dose after 5-10 minutes for a maximum of two doses. Contact 911 and monitor vital signs.  
 Other: \_\_\_\_\_

**9. NURSING ORDERS**

Skilled nursing to insert, maintain and remove/de-access vascular access daily, weekly and/or as needed, draw labs as ordered, conduct patient assessments, and educate patient/caregiver on home infusion, medication administration, self-monitoring, and patient safety.  
Skilled Nursing Services Needed?  Yes  No Additional Instructions: \_\_\_\_\_

**10.**  Dispense as written  Substitution Permitted

**PHYSICIAN'S SIGNATURE (required):** \_\_\_\_\_ **Date:** \_\_\_\_\_